

PATTERNS OF variation in small mammal abundance in the Road Corridor, 1992- 2000




Eric Rexstad and Ed Debevec

Institute of Arctic Biology
University of Alaska Fairbanks




Utility of microtines for ecological monitoring




Challenges of monitoring small mammals in Denali National Park and Preserve






Vertebrate data geyser




- Two paid technicians have, on average, had >2400 encounters per year
- Totaling 22,000+ encounters during the project


Small mammal abundance estimates for RR2: 1992-2000




Small mammal abundance estimates for RF1: 1992-2000



Intra-annual temporal variation in Clethrionomys dynamics




Small mammal abundance estimates for RF1: 1992-2000



Rock Cr.

Small mammal abundance estimates for W1: 1995-2000



McKinley
Bar

- Contrasting similar habitats at opposite ends of the Road Corridor,
- *Clethrionomys* appears to have similar temporal patterns,
- *Microtus*, however exhibit more temporal variation at the eastern end of the road than at the western end

Unraveling the temporal variability in microtine abundance

$$\begin{aligned}\log[E(N)] = & \beta_0 + \beta_1 W + \beta_2 C + \beta_3 R + \beta_4 WSI + \beta_5 SOI + \beta_6 SRI \\ & + \beta_7 C * R + \beta_8 C * WSI + \beta_9 + C * SOI + \beta_{10} C * SRI \\ & + \beta_{11} R * WSI + \beta_{12} R * SOI + \beta_{13} R * SRI \\ & + \beta_{14} C * R * WSI + \beta_{15} C * R * SOI + \beta_{16} C * R * SRI\end{aligned}$$

where

N = end-of-summer vole abundance,

W = 1 if web and 0 if grid,

C = 1 if *Clethrionomys* and 0 if *Microtus*,

R = 1 if riparian and 0 if forest,


WSI = winter severity index,

SOI = spring onset index, and


SRI = spring rainfall index.

In fitting the model, we used quasi-likelihood methods to produce variance estimates appropriate for data with probable extra-Poisson variation. Because our response variable was itself an estimate with a measure of uncertainty, we weighted each observation by the inverse of its coefficient of variation.

Clethrionomys - Forest



Clethrionomys - Riparian






Model performance

- Our abiotic predictive model “absorbs” variation in microtine abundance attributable to annual variation in climatic factors.
- Remaining variation may be used to detect signs of “stress” in these populations

Spatial variation in microtine abundance

- At our intermediate level of spatial sampling (1997-1999), we found no evidence of spatial variation in microtine abundance between watersheds situated within ~50km² region directly west of Denali Park headquarters.
- We are still assessing spatial variation from the 2000 field season





Track continued progress at:

<http://mercury.bio.uaf.edu/~erexstad.faculty/rexst.html>

